



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 498/16, 495/04, 487/04, 471/04, 498/04, A61K 31/55 // (C07D 498/16, 333:00, 261:00, 223:00) (C07D 498/16, 261:00, 243:00, 209:00) (C07D 495/04, 333:00, 223:00) (C07D 487/04, 243:00, 209:00) (C07D 471/04, 223:00, 221:00) (C07D 498/04, 267:00, 221:00)</p>	<p>A1</p>	<p>(11) International Publication Number: WO 97/39001</p> <p>(43) International Publication Date: 23 October 1997 (23.10.97)</p>
<p>(21) International Application Number: PCT/EP97/01830</p> <p>(22) International Filing Date: 9 April 1997 (09.04.97)</p> <p>(30) Priority Data: 96200991.6 12 April 1996 (12.04.96) EP</p> <p>(34) Countries for which the regional or international application was filed: DE et al.</p> <p>(71) Applicant (for all designated States except US): JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): ANDRÉS-GIL, José Ignacio [ES/ES]; Vesubiana, 22, Getafe-Sector 3, E-28905 Madrid (ES). MARTINEZ, Pedro [ES/ES]; Radio 28, 22A, E-28109 Madrid (ES). FERNÁNDEZ-GADEA, Francisco Javier [ES/ES]; Sagra, 20, Cerro De Las Perdices, E-45593 Toledo (ES). SIPIDO, Victor, Karel [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE).</p>		<p>(74) Agent: QUAGHEBEUR, Luc; Janssen Pharmaceutica N.V., Patent Dept., Turnhoutseweg 30, B-2340 Beerse (BE).</p> <p>(81) Designated States: AL, AM, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KR, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>
<p>(54) Title: ISOXAZOLIDINE DERIVATIVES</p>		
<p>(57) Abstract</p> <p>This invention concerns the compounds of formula (I), the <i>N</i>-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein <i>n</i> is zero to 6; <i>p</i> is zero to 3; <i>q</i> is zero to 3; <i>r</i> is zero to 3; <i>R</i>¹ and <i>R</i>² each independently are hydrogen; optionally substituted C₁₋₆alkyl; C₁₋₆alkylcarbonyl; trihalomethylcarbonyl; or <i>R</i>¹ and <i>R</i>² taken together with the nitrogen atom to which they are attached may form a morpholinyl ring or an optionally substituted heterocycle; each <i>R</i>³ and <i>R</i>⁴ independently are halo, cyano, hydroxy, trihalomethyl, trihalomethoxy, carboxyl, nitro, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylcarbonylamino, aminosulfonyl, mono- or di(C₁₋₆alkyl)-aminosulfonyl, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylcarbonyl or C₁₋₆alkyloxycarbonyl; each <i>R</i>⁵ independently is C₁₋₆alkyl, cyano or trihalomethyl; <i>X</i> is CR⁶R⁷, NR⁸, O, S, S(=O) or S(=O)₂; each formula (i) independently represents a bivalent aromatic heterocycle wherein the heterocycle may be selected from the group consisting of pyrrole, pyrazole, imidazole, triazole, furane, thiophene, isoxazole, oxazole, isothiazole, thiazole, pyridine, pyridazine, pyrimidine, pyrazine and triazine; or one of the two bivalent aromatic heterocycles in the compounds of formula (I) may be 1,2-benzenediyl; and aryl is optionally substituted phenyl; it further relates to compositions comprising these compounds, as well as their use as therapeutic agents in the treatment or the prevention of CNS disorders, cardiovascular disorders or gastrointestinal disorders.</p> <div style="text-align: center;"> <p>(I)</p> </div> <div style="text-align: center;"> <p>(II)</p> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-


ISOXAZOLIDINE DERIVATIVES

This invention concerns substituted tetracyclic isoxazolidine derivatives having antipsychotic, cardiovascular and gastrokinetic activity and their preparations; it further relates to compositions comprising them, as well as their use as a medicine.

US 4,039,558 discloses pyrrolidinodibenzo-azepine, -oxazepine, -thiazepine and -diazepine derivatives having antihistamine, sedative and antidepressant properties.

EP-A-0,421,823 describes dibenzopyrazino- or benzo-pyrido-pyrazino-azepine derivatives having anti-allergic and anti-asthmatic activities. The present compounds differ therefrom by the presence of an isoxazolidine ring, and by their pharmacological properties.

This invention concerns compounds of formula (I)



the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein :

n is zero, 1, 2, 3, 4, 5, or 6;

p is zero, 1, 2 or 3;


q is zero, 1, 2 or 3;

r is zero, 1, 2 or 3;


*R*¹ and *R*² each independently are hydrogen; C₁₋₆alkyl; C₁₋₆alkylcarbonyl;

trihalomethylcarbonyl; C₁₋₆alkyl substituted with hydroxy, C₁₋₆alkyloxy, carboxyl,


C₁₋₆alkylcarbonyloxy, C₁₋₆alkyloxycarbonyl or aryl; or *R*¹ and *R*² taken together with the nitrogen atom to which they are attached may form a morpholinyl ring or a radical of formula :




(a)



(b)



(c)



wherein :

R^9 , R^{10} , R^{11} and R^{12} each independently are hydrogen, halo, trihalomethyl, or C_{1-6} alkyl;

5 m is zero, 1, 2, or 3;

R^{13} , R^{14} , R^{15} and R^{16} each independently are hydrogen or C_{1-6} alkyl; or

R^{15} and R^{16} taken together may form a bivalent radical C_{4-5} alkanediyl;

R^{17} is hydrogen; C_{1-6} alkyl; C_{1-6} alkylcarbonyl; trihalomethylcarbonyl;

10 C_{1-6} alkyloxycarbonyl; aryl; di(aryl)methyl; C_{1-6} alkyl substituted with hydroxy, C_{1-6} alkyloxy, carboxyl, C_{1-6} alkylcarbonyloxy, C_{1-6} alkyloxycarbonyl or aryl;

each R^3 independently is halo, cyano, hydroxy, trihalomethyl, trihalomethoxy, carboxyl, nitro, amino, mono- or di(C_{1-6} alkyl)amino, C_{1-6} alkylcarbonylamino, aminosulfonyl, mono- or di(C_{1-6} alkyl)aminosulfonyl, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{1-6} alkylcarbonyl or C_{1-6} alkyloxycarbonyl;

15 each R^4 independently is halo, cyano, hydroxy, trihalomethyl, trihalomethoxy, carboxyl, nitro, amino, mono- or di(C_{1-6} alkyl)amino, C_{1-6} alkylcarbonylamino, aminosulfonyl, mono- or di(C_{1-6} alkyl)aminosulfonyl, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{1-6} alkylcarbonyl or C_{1-6} alkyloxycarbonyl;

each R^5 independently is C_{1-6} alkyl, cyano or trihalomethyl;


20 X is CR^6R^7 , NR^8 , O, S, $S(=O)$ or $S(=O)_2$; wherein

R^6 and R^7 each independently are hydrogen, hydroxy, C_{1-6} alkyl, trihalomethyl, C_{1-6} alkyloxy or R^6 and R^7 taken together may form methylene; mono- or

di(cyano)methylene; a bivalent radical of formula $-(CH_2)_2-$, $-(CH_2)_3-$, $-(CH_2)_4-$, $-(CH_2)_5-$, $-O-(CH_2)_2-O-$, $-O-(CH_2)_3-O-$; or, together with the carbon atom to which


25 they are attached, a carbonyl;

R^8 is hydrogen, C_{1-6} alkyl, C_{1-6} alkylcarbonyl, arylcarbonyl, aryl C_{1-6} alkylcarbonyl, C_{1-6} alkylsulfonyl, arylsulfonyl or aryl C_{1-6} alkylsulfonyl;

each  independently represents a bivalent aromatic heterocycle wherein the

heterocycle may be selected from the group consisting of pyrrole, pyrazole, imidazole, 30 triazole, furane, thiophene, isoxazole, oxazole, isothiazole, thiazole, pyridine,

pyridazine, pyrimidine, pyrazine and triazine; or one of the two bivalent aromatic heterocycles in the compounds of formula (I) may be 1,2-benzenediyl; and aryl is phenyl; or phenyl substituted with 1, 2 or 3 substituents selected from halo, hydroxy, C₁₋₆alkyl and trihalomethyl.

5 In the foregoing definitions C₁₋₆alkyl defines straight and branch chained saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as, for example, methyl, ethyl, propyl, butyl, 1-methylpropyl, 1,1-dimethylethyl, pentyl, hexyl; C₄₋₅alkanediyl defines bivalent straight and branch chained saturated hydrocarbon
10 radicals having from 4 to 5 carbon atoms such as, for example, 1,4-butanediyl, 1,5-pentanediyl; halo is generic to fluoro, chloro, bromo and iodo. The term monocyanomethylene stands for a radical of formula =CHCN, and dicyanomethylene for a radical of formula =C(CN)₂. The term halomethyl is meant to include mono-, di-, and trihalomethyl. Examples of halomethyl are fluoromethyl, difluoromethyl and
15 particularly trifluoromethyl. In case R⁶ and R⁷ taken together form a bivalent radical of formula -(CH₂)₂-, -(CH₂)₃-, -(CH₂)₄-, -(CH₂)₅-, -O-(CH₂)₂-O- or -O-(CH₂)₃-O-, the compounds of formula (I) are spiro compounds. The bivalent heterocyclic radicals in the definition of  include all bivalent radicals wherein the radical positions are adjacent on the heterocycle.

20 The pharmaceutically acceptable addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic base and acid addition salt forms which the compounds of formula (I) are able to form. The acid addition salt form of a compound of formula (I) that occurs in its free form as a base can be obtained by
25 treating the base form of the compound of formula (I) with an appropriate acid such as an inorganic acid, for example, hydrohalic acid, e.g. hydrochloric or hydrobromic, sulfuric, nitric, phosphoric and the like acids; or an organic acid, such as, for example, acetic, hydroxyacetic, propanoic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic,
30 *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

The compounds of formula (I) containing acidic protons may be converted into their therapeutically active non-toxic base, *i.e.* metal or amine. addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms
35 comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with

organic bases, e.g. the benzathine, *N*-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

5 Conversely said salt forms can be converted into the free forms by treatment with an appropriate base or acid.

10 The term addition salt as used hereinabove also comprises the solvates which the compounds of formula (I) as well as the salts thereof, are able to form. Such solvates are for example hydrates, alcoholates and the like.

15 The *N*-oxide forms of the compounds of formula (I) are meant to comprise those compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide, particularly those *N*-oxides wherein the nitrogen bearing the R¹ and R² substituents is *N*-oxidized.

20 The term "stereochemically isomeric forms" as used hereinbefore and hereinafter defines all the possible isomeric forms in which the compounds of formula (I) may occur. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture, and in particular the racemic mixture, of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure. Stereochemically isomeric forms of the compounds of formula (I) and mixtures of such forms are obviously intended to be encompassed by formula (I).


25 The numbering of the isoxazolidine ring-system present in the compounds of formula (I), is used hereinafter as defined by Chemical Abstracts nomenclature. The compounds of formula (I) occur as *cis* and *trans* isomers. Said terms refer to the position of the substituents on the isoxazolidine ring and are also in accordance with Chemical Abstracts nomenclature. The nomenclature is unusual in that the only relevant substituent of
30 carbon atom 3a is either hydrogen or R⁵. When establishing the configuration, the substituent on carbon atom 3a and the substituent with the highest priority on carbon atom 2 (*i.e.* either $-(CH_2)_n-NR^1R^2$ or R⁵) are considered. When the relevant substituent on carbon atom 3a and the substituent with the highest priority on carbon atom 2 are on the same side of the mean plane determined by the isoxazolidine ring then the
35 configuration is designated *cis*, if not, the configuration is designated *trans*.

The compounds of formula (I) have at least two asymmetric centers, namely carbon atom 3a and carbon atom 2. Said asymmetric centers and any other asymmetric center which may be present, are indicated by the descriptors R and S. When a


monocyanomethylene moiety is present in the compounds of formula (I), said moiety may have the E- or Z-configuration.


- Whenever used hereinafter, the term "compounds of formula (I)" is meant to also include the pharmaceutically acceptable acid addition salts, base addition salts and all stereoisomeric forms, and also the *N*-oxide forms.

Interesting compounds are those compounds of formula (I) wherein one of the two



radicals is 1,2-benzenediyl; in particular those compounds of formula (I)

- wherein one of the two  radicals is 1,2-benzenediyl and the other bivalent

radical  is a bivalent pyridine, imidazole, pyrrole or thiophene radical.

- Other interesting compounds are those compounds of formula (I) wherein X is CR⁶R⁷ or O.

Still other interesting compounds are those compounds of formula (I) wherein r is zero.

- Preferred compounds are those compounds of formula (I) wherein R¹ and R² are both methyl and n is 1 or 2.


Other preferred compounds are those compounds of formula (I) wherein R¹ and R² are taken together with the nitrogen atom to which they are attached to form a morpholinyl ring or a radical of formula (e).

- Also preferred are those compounds of formula (I) wherein R¹ is hydrogen or C₁₋₆alkyl substituted with C₁₋₆alkylcarbonyloxy, R² is methyl and n is 1 or 2.


Most preferred compounds are :

- 2,3,3a,8-tetrahydro-*N,N*-dimethylisoxazolo[2,3-a]pyrrolo[2,1-c][1,4]benzo-diazepine-2-methanamine;
 2,3,3a,8-tetrahydro-*N,N*-dimethylimidazo[2,1-c]isoxazolo[2,3-a][1,4]benzo-diazepine-2-methanamine;
 2,3,3a,7-tetrahydro-*N,N*-dimethylisoxazolo[2,3-a]thieno[2,3-c][1]benz-azepine-2-methanamine;

2,3,3a,7-tetrahydro-*N,N*-dimethylisoxazolo[2,3-*a*]thieno[3,2-*c*][1]benz-azepine-2-methanamine; the stereochemically isomeric forms and pharmaceutically acceptable addition salts thereof, and also the *N*-oxide forms thereof.

- 5 In general, the compounds of formula (I) may be prepared by a 1,3-dipolar cycloaddition of a dienophile of formula (III) and an intermediate of formula (II). In the intermediates (II) and (III) and in any other intermediate mentioned hereinunder, R^1 to R^5 , X , n , r , p , q and  are as defined hereinabove, unless otherwise indicated.

- Further, the hydrogens on the sp^2 -hybridized aliphatic carbon atoms in the intermediates (II) and (III) may each individually be replaced by R^5 . Said 1,3-dipolar cycloaddition may conveniently be carried out by mixing the reactants, optionally in a reaction-inert solvent such as, for example, toluene, 4-methyl-2-pentanone or tetrahydrofuran, or a mixture of such solvents. Stirring and elevated temperatures, or increased pressure may enhance the rate of the reaction. The reaction of intermediate (II) with intermediate (III) in practice is regioselective yielding compounds of formula (I).




- In this and the following preparations, the reaction products may be isolated from the reaction medium and, if necessary, further purified according to methodologies generally known in the art such as, for example, extraction, crystallization, trituration and chromatography.

- The compounds of formula (I) may also be converted into each other following art-known transformation reactions. For instance,
- 25 a) a compound of formula (I), wherein R^1 and R^2 taken together with the nitrogen atom to which they are attached form a radical of formula (b), may be converted into the corresponding primary amine by treatment with hydrazine or aqueous alkali;
 - b) a compound of formula (I), wherein R^1 or R^2 is trifluoromethylcarbonyl, may be converted into the corresponding primary or secondary amine by hydrolysis with
 - 30 aqueous alkali;

- c) a compound of formula (I), wherein R^1 or R^2 is C_{1-6} alkyl substituted with C_{1-6} alkylcarbonyloxy may be hydrolyzed into a compound of formula (I) wherein R^1 or R^2 is C_{1-6} alkyl substituted with hydroxy;
- d) a compound of formula (I), wherein R^1 and R^2 are both hydrogen may be mono- or di-*N*-alkylated to the corresponding amine form;
- e) a compound of formula (I), wherein R^1 and R^2 are both hydrogen may be *N*-acylated to the corresponding amide;
- f) a compound of formula (I), containing a C_{1-6} alkyloxycarbonyl group may be hydrolyzed to the corresponding carboxylic acid.


In addition, the compounds of formula (I) wherein X is other than S may also be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with 3-phenyl-2-(phenylsulfonyl)oxaziridine or with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chlorobenzenecarboperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. *t*-butyl hydroperoxide. Suitable solvents are, for example, water, lower alkanols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.


The intermediates of formula (II) wherein X^a is CR^6R^7 , O, $S(=O)$ or $S(=O)_2$, said intermediates are represented by formula (II-a), may be prepared by the oxidation of an intermediate of formula (IV) with a suitable oxidizing agent such as, for example, 2-benzenesulfonyl-3-phenyl-oxaziridine, hydrogen peroxide, *tert*-butyl hydroxyperoxide, or metachloroperbenzoic acid.




Said oxidation is performed in a reaction-inert solvent at temperatures ranging between -20°C and 50°C , preferably between 0°C and room temperature. Suitable solvents are,

for example, water, dichloromethane, toluene, methanol, 4-methyl-2-pentanone, or a mixture of such solvents. When using peroxide oxidants, the reaction rate may be enhanced by using metallic catalysts such as, for example, Na_2WO_4 , $\text{VO}(\text{acetylacetonate})_2$, $\text{Ti}(\text{OBu})_4$, or $\text{MoO}_2(\text{acetylacetonate})_2$, optionally under a
 5 reaction-inert atmosphere such as, for example, argon.

Alternatively, intermediates of formula (II) wherein R^{3b} is the same as R^3 but other than nitro, and R^{4b} is the same as R^4 but other than nitro, said intermediates are represented by formula (II-b), may be prepared by reducing the nitro group of an
 10 intermediate of formula (V) in the presence of water and a suitable reducing agent such as, for example, zinc or iron; subsequently followed by an *in situ* intramolecular cyclization in the presence of a weak acid such as, for example, ammoniumchloride or acetic acid. Said reductive cyclization is performed in a reaction-inert solvent such as, for example, 1,4-dioxane. Stirring and elevated temperatures may enhance the rate of
 15 the reaction. In intermediate (V), R^{3b} and R^{4b} are defined as in intermediates of formula (II-b) and both the formyl and nitro substituent are on an atom adjacent to the -X radical on the respective aromatic heterocycles of formula  as defined in the compounds of formula (I).



20 The intermediates of formula (II) wherein , X, R^3 , R^4 , p and q are as defined hereinabove are deemed novel.

Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures. Diastereomers may be separated by
 25 physical methods such as selective crystallization and chromatographic techniques, e.g. counter-current distribution, liquid chromatography and the like.

The compounds of formula (I) as prepared in the hereinabove described processes are generally racemic mixtures of enantiomers which can be separated from one another
 30 following art-known resolution procedures. The racemic compounds of formula (I) which are sufficiently basic or acidic may be converted into the corresponding

diastereomeric salt forms by reaction with a suitable chiral acid respectively with a suitable chiral base. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali or acid. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography using a chiral stationary phase. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

The compounds of the present invention show affinity for 5-HT₂ receptors, particularly for 5-HT_{2A} and 5-HT_{2C} receptors (nomenclature as described by D. Hoyer in "Serotonin (5-HT) in neurologic and psychiatric disorders" edited by M.D. Ferrari and published in 1994 by the Boerhaave Commission of the University of Leiden). The serotonin antagonistic properties of the present compounds may be demonstrated by their inhibitory effect in the "5-hydroxytryptophan Test on Rats" which is described in Drug Dev. Res., 13, 237-244 (1988). Furthermore, the present compounds show interesting pharmacological activity *in vivo* in the "mCPP Test on Rats" which is described hereinafter, in the "Elevated and Illuminated Plus Maze Test" which is described in Drug Dev. Res., 18, 119-144 (1989), and in the "Combined Apomorphine, Tryptamine, Norepinephrine (ATN) Test on Rats" which is described in Arch. Int. Pharmacodyn, 227, 238-253 (1977).

In view of these pharmacological and physicochemical properties, the compounds of formula (I) are useful as therapeutic agents in the treatment or the prevention of central nervous system disorders like anxiety, depression and mild depression, bipolar disorders, sleep- and sexual disorders, psychosis, borderline psychosis, schizophrenia, migraine, personality disorders or obsessive-compulsive disorders, social phobias or panic attacks, organic mental disorders, mental disorders in children, aggression, memory disorders and attitude disorders in older people, addiction, obesity, bulimia and similar disorders. In particular, the present compounds may be used as anxiolytics, antipsychotics, antidepressants, anti-migraine agents and as agents having the potential to overrule the addictive properties of drugs of abuse.

The compounds of formula (I) may also be used as therapeutic agents in the treatment of motoric disorders. It may be advantageous to use the present compounds in combination with classical therapeutic agents for such disorders.

- 5 The compounds of formula (I) may also serve in the treatment or the prevention of damage to the nervous system caused by trauma, stroke, neurodegenerative illnesses and the like; cardiovascular disorders like high blood pressure, thrombosis, stroke, and the like; and gastrointestinal disorders like dysfunction of the motility of the gastrointestinal system and the like.

- 10 In view of the above uses of the compounds of formula (I), it follows that the present invention also provides a method of treating warm-blooded animals suffering from such diseases, said method comprising the systemic administration of a therapeutic amount of a compound of formula (I) effective in treating the above described disorders, in particular, in treating anxiety, psychosis, depression, migraine and
15 addictive properties of drugs of abuse.

- The present invention thus also relates to compounds of formula (I) as defined hereinabove for use as a medicine, in particular, the compounds of formula (I) may be
20 used for the manufacture of a medicine for treating anxiety, psychosis, depression, migraine and addictive properties of drugs of abuse.

- Those of skill in the treatment of such diseases could determine the effective therapeutic daily amount from the test results presented hereinafter. An effective
25 therapeutic daily amount would be from about 0.01 mg/kg to about 10 mg/kg body weight, more preferably from about 0.05 mg/kg to about 1 mg/kg body weight.

- For ease of administration, the subject compounds may be formulated into various pharmaceutical forms for administration purposes. To prepare the pharmaceutical
30 compositions of this invention, a therapeutically effective amount of the particular compound, optionally in addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable,
35 preferably, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions,

5 syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable solutions containing compounds of formula (I) may be formulated in an oil for prolonged action. Appropriate oils for this purpose are, for example, peanut oil, sesame oil, cottonseed oil, corn oil, soy bean oil, synthetic glycerol esters of long chain fatty acids and mixtures of these and other oils. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wettable agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause any significant deleterious effects on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on or as an ointment. Acid or base addition salts of compounds of formula (I) due to their increased water solubility over the corresponding base or acid form, are obviously more suitable in the preparation of aqueous compositions.

25 In order to enhance the solubility and/or the stability of the compounds of formula (I) in pharmaceutical compositions, it can be advantageous to employ α -, β - or γ -cyclodextrins or their derivatives, in particular hydroxyalkyl substituted cyclodextrins, e.g. 2-hydroxypropyl- β -cyclodextrin. Also co-solvents such as alcohols may improve the solubility and/or the stability of the compounds of formula (I) in pharmaceutical compositions.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers,

injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

5 The following examples are intended to illustrate and not to limit the scope of the present invention.

Experimental part

A. Preparation of the intermediate compounds

Example A.1

- 10 a) Sodium hydride (60%) (0.0899 mol) was added portionwise under N₂ at room temperature to a mixture of 2-hydroxybenzaldehyde (0.0818 mol) in *N,N*-dimethylformamide and the mixture was stirred under N₂ for 15 minutes. 2-Chloro-3-nitropyridine (0.0818 mol) in *N,N*-dimethylformamide was added and the mixture was stirred at 100°C for 2 hours. The mixture was poured into ice/water, extracted with
15 diethylether and the solvent was evaporated, yielding 4.5 g (23%) of 2-[(3-nitro-2-pyridinyl)oxy]benzaldehyde (interm. 1).
- b) Zinc (0.0821 mol) was added to a suspension of intermediate (1) (0.0158 mol) in 1,4-dioxane (100 ml) and water (6.2 ml). Ammonium chloride (0.0413 mol) was added dropwise at 5-7°C and the mixture was stirred at 5°C for 1 hour. The precipitate was
20 filtered off and the filtrate was evaporated. The residue (6.6 g) was treated with water, extracted with CH₂Cl₂ and the solvent was evaporated. The product was used without further purification, yielding 3.7 g of pyrido[2,3-*b*][1,4]benzoxazepine, 5-oxide (interm. 2).


Example A.2

- 25 a) A mixture of 2-aminobenzyl alcohol (0.1218 mol), 3-thiophenecarboxaldehyde (0.1218 mol) and molecular sieves (3 Å) in 2-propanol (150 ml) was stirred overnight at room temperature. The crude reaction mixture was filtered, the residue was treated with CH₂Cl₂, and the mixture was filtered again. The solvent was evaporated, yielding 21.53 g (81%) of (±)-1,4-dihydro-2-(3-thienyl)-2*H*-3,1-benzoxazine (interm. 3).
- 30 b) Sodium tetrahydroborate (0.1979 mol) was added portionwise to intermediate (3) (0.0989 mol) in ethanol (230 ml), stirred at room temperature. The reaction mixture was stirred for 30 minutes at room temperature, then for 90 minutes at reflux temperature. The solvent was evaporated, the residue was treated with a saturated aqueous NH₄Cl solution, and the resulting mixture was extracted with CH₂Cl₂. The
35 separated organic layer was dried, filtered and the solvent evaporated, yielding 20.89 g (85%) of 2-[(3-thienylmethyl)amino]benzenemethanol (interm. 4).

- c) Tin (IV) chloride (0.2658 mol) was added dropwise to intermediate (4) (0.088 mol) in 1,2-dichloroethane (1000 ml). The reaction mixture was stirred for one hour at 80 °C. The reaction mixture was cooled on an ice bath. Water was added dropwise and the resulting solution was basified, then extracted with CH₂Cl₂. The separated organic layer was dried, filtered and the solvent evaporated. The residue was purified by flash column chromatography over silica gel (eluent: hexane/ethylacetate 9/1 and 8/2). The desired fractions were collected and the solvent was evaporated, yielding 4.76 g (27%) of 5,10-dihydro-4*H*-thieno[3,2-*c*][1]benzazepine (interm. 5).
- d) Intermediate (5) (0.0145 mol) was dissolved in CH₂Cl₂ (128 ml) and cooled on an ice bath. (±)-3-Phenyl-2-(phenylsulfonyl)oxaziridine (0.02911 mol) was added portionwise and the reaction mixture was stirred for 4 hours. The solvent was evaporated and the residue was purified by open column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 100/0 and 96/4). The pure fractions were collected and the solvent was evaporated, yielding 3.64 g (82%) 10*H*-thieno[3,2-*c*][1]benzazepine, 5-oxide (interm. 6).


Tables 1 and 2 list the intermediates of formula (II) which were made according to one of the above examples (column Ex. No.).

Table 1



Interm. No.	Ex. No.	Ar
6	A.2	
7	A.1	
8	A.1	
9	A.2	
10	A.2	

Table 2



Interm. No.	Ex. No.	X	Ar
2	A.1	O	
11	A.2	CH ₂	

B. Preparation of the compounds of formula (I)

5 Example B.1

A mixture of intermediate (2) (0.0174 mol) and 3-amino-1-propene (0.0261 mol) in tetrahydrofuran (60 ml) was stirred at 60°C for 5 hour. The solvent was evaporated and the residue was purified by flash chromatography over silica gel (eluent : CH₂Cl₂/hexane/2-propapnone 6/3/1 and CH₂Cl₂/hexane/CH₃OH 6/3.5/0.5). The pure fractions were collected and the solvent evaporated. The residue (2.5g) was washed with diethylether and purified further by HPLC (eluent : hexane/CH₂Cl₂/CH₃OH 6/3/1). The pure fractions were collected and evaporated, yielding 1.4 g (27%) of (±)-*cis*-3,13b-dihydro-*N,N*-dimethyl-2*H*-isoxazolo[2,3-*d*]pyrido[2,3-*b*][1,4]benzoxazepine-2-methanamine (comp. 1; mp. 106.3°C).

15 Example B.2

(±)-*cis*-2,3,3a,7-tetrahydro-*N,N*-dimethylisoxazolo[2,3-*a*]thieno[3,2-*c*][1]benzazepine-2-methanamine ethanedioate(1:1) (comp. 2; mp. 136.9°C) was prepared in a similar way as compound 1 in example B.1 but using toluene as a solvent.

Example B.3

20 (±)-*cis*-2,3,3a,8-tetrahydro-*N,N*-dimethylisoxazolo[2,3-*a*]pyrrolo[2,1-*c*][1,4]-benzodiazepine-2-methanamine ethanedioate (1:1) (comp. 3; mp. 137.6°C) was prepared in a similar way as compound 1 in example B.1 but using a mixture of tetrahydrofuran and toluene as a solvent.


Example B.4

25 2-(methyl-2-propenylamino)ethanol acetate(ester) (0.0176 mol) was added to a solution of intermediate (11) (0.0160 mol) in tetrahydrofuran (110 ml) and the resulting reaction mixture was stirred and refluxed for 24 hours. Toluene (100 ml) was added and

the mixture was stirred and refluxed for 24 hours. The solvent was evaporated and the residue was purified by short open column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$ 96/4), then by HPLC (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 96/4). The pure fractions were collected and the solvent was evaporated, yielding 0.360 g (10%) of (\pm)-
 5 2,3,3a,8-tetrahydro-2-(4-morpholinylmethyl)isoxazolo[3,2-a]pyrido[3,4-c][2]-benzazepine (comp. 9; mp. 46°C).


Tables 3 and 4 list the compounds of formula (I) which were made according to one of the above examples (column "Ex. No.").

10 Table 3



Co. No	Ex. No.	Ar	physical data
2	B.2		(\pm)- <i>cis</i> ; mp. 136.9°C; oxalic acid (1:1)
3	B.3		(\pm)- <i>cis</i> ; mp. 137.6°C; oxalic acid (1:1)
4	B.1		(\pm)- <i>cis</i> ; mp. 135.5°C; H_2O (1:2).HCl (1:2)
5	B.2		(\pm)- <i>cis</i> ; mp. 157.8°C; oxalic acid (1:1)
6	B.1		mp. 112.5°C; oxalic acid (1:1). H_2O (1:1) .2-propanolate (1:1)

Table 4



Co. No.	Ex. No.	X	R	Ar	Physical data
1	B.1	O	$-N(CH_3)_2$		(±)- <i>cis</i> ; mp. 106.3°C
7	B.1	CH_2	$-N(CH_3)_2$		(±)- <i>cis</i>
8	B.1	CH_2			(±)- <i>cis</i>
9	B.4	CH_2			(±)- <i>cis</i> ; mp. 46°C
10	B.1	CH_2			(±)- <i>cis</i> ; mp. 116°C

C. Pharmacological example

5 Example C.1 : "mCPP Test on Rats"

Rats were treated with the test compound at a dose varying between 0.0025 mg/kg and 40 mg/kg body weight at pre-test time T of 1 hour, and with 1 mg/kg mCPP (metachlorophenylpiperazine), injected intravenously, 15 minutes prior to the test.

- 10 After pre-test time T elapsed, treated rats were submitted to the "Open Field Test on Rats" as described in Drug Dev. Res. 18, 119-144 (1989), but using an infra-red light source instead of a Kleverlux® (12V/20W) light source. A dose at which 42 % of the tested rats showed suppression of the mCPP induced effects, *i.e.* mCPP-antagonism, was defined as an active dose. Compound numbers 2, 3 and 7 were active at a test dose of 10 mg/kg or less.

15 Example C.2: "Elevated and Illuminated Plus Maze Test on Rats"

The "Elevated and Illuminated Plus Maze Test on Rats" is described in Drug Dev. Res. 18, 119-144 (1989). An active dose of a test compound in said test was defined as a dose at which 42 % of the tested rats explored the illuminated arms of the maze. The compounds numbers 2, 4 and 5 showed activity at a test dose of 2.5 mg/kg or less.

Example C.3 : *In vitro* binding affinity for 5-HT_{2A} and 5-HT_{2C} receptors

The interaction of the compounds of formula (I) with 5-HT_{2A} and 5-HT_{2C} receptors was assessed in *in vitro* radioligand binding experiments.

- 5 In general, a low concentration of a radioligand with a high binding affinity for the receptor is incubated with a sample of a tissue preparation enriched in a particular receptor (1 to 5 mg tissue) in a buffered medium (0.2 to 5 ml). During the incubation, the radioligands bind to the receptor. When equilibrium of binding is reached, the receptor bound radioactivity is separated from the non-bound radioactivity, and the
- 10 receptor bound activity is counted. The interaction of the test compounds with the receptors is assessed in competition binding experiments. Various concentrations of the test compound are added to the incubation mixture containing the tissue preparation and the radioligand. Binding of the radioligand will be inhibited by the test compound in proportion to its binding affinity and its concentration.
- 15 The radioligand used for 5-HT_{2A} binding affinity is ³H-ketanserin and the tissue used is the frontal cortex of the rat. At a test concentration of 10⁻⁷ M, the compounds with number 2, 3, 5, and 7 produced an inhibition of the 5-HT_{2A} receptor of more than 40 %, and the other compounds produced an inhibition of less than 40 %.
- 20 The radioligand used for 5-HT_{2C} binding affinity is ³H-mesulergine and the tissue used is the choroid plexus of the pig. At a test concentration of 10⁻⁷ M, the compounds with number 2, 3, 5, 6, and 7 produced an inhibition of the 5-HT_{2C} receptor of more than 40 %, and the other compounds produced an inhibition of less than 40 %.

D. Composition examples

- 25 "Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I), a pharmaceutically acceptable acid addition salt, a stereochemically isomeric form thereof or a *N*-oxide form thereof.

Example D.1 : ORAL SOLUTION

- Methyl 4-hydroxybenzoate (9 g) and propyl 4-hydroxybenzoate (1 g) were dissolved in boiling purified water (4 l). In 3 l of this solution were dissolved first
- 30 2,3-dihydroxybutanedioic acid (10 g) and thereafter A.I (20 g). The latter solution was combined with the remaining part of the former solution and 1,2,3-propanetriol (12 l) and sorbitol 70% solution (3 l) were added thereto. Sodium saccharin (40 g) were dissolved in water (500 ml) and raspberry (2 ml) and gooseberry essence (2 ml) were added. The latter solution was combined with the former, water was added q.s. to a
- 35 volume of 20 l providing an oral solution comprising 5 mg of the active ingredient per teaspoonful (5 ml). The resulting solution was filled in suitable containers.

Example D.2 : FILM-COATED TABLETSPreparation of tablet core

A mixture of A.I. (100 g), lactose (570 g) and starch (200 g) was mixed well and thereafter humidified with a solution of sodium dodecyl sulfate (5 g) and
5 polyvinylpyrrolidone (10 g) in water (200 ml). The wet powder mixture was sieved, dried and sieved again. Then there was added microcrystalline cellulose (100 g) and hydrogenated vegetable oil (15 g). The whole was mixed well and compressed into tablets, giving 10.000 tablets, each containing 10 mg of the active ingredient.

Coating


10 To a solution of methyl cellulose (10 g) in denaturated ethanol (75 ml) there was added a solution of ethyl cellulose (5 g) in dichloromethane (150 ml). Then there were added dichloromethane (75 ml) and 1,2,3-propanetriol (2.5 ml). Polyethylene glycol (10 g) was molten and dissolved in dichloromethane (75 ml). The latter solution was added to the former and then there were added magnesium octadecanoate (2.5 g),
15 polyvinylpyrrolidone (5 g) and concentrated colour suspension (30 ml) and the whole was homogenated. The tablet cores were coated with the thus obtained mixture in a coating apparatus.

Example D.3 : INJECTABLE SOLUTION

Methyl 4-hydroxybenzoate (1.8 g) and propyl 4-hydroxybenzoate (0.2 g) were
20 dissolved in boiling water (500 ml) for injection. After cooling to about 50°C there were added while stirring lactic acid (4 g), propylene glycol (0.05 g) and A.I. (4 g). The solution was cooled to room temperature and supplemented with water for injection q.s. ad 1 l, giving a solution comprising 4 mg/ml of A.I.. The solution was sterilized by filtration and filled in sterile containers.

Claims

1. A compound of formula



5 a *N*-oxide form, a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof, wherein :




n is zero, 1, 2, 3, 4, 5, or 6;

p is zero, 1, 2 or 3;


q is zero, 1, 2 or 3;

r is zero, 1, 2 or 3;


10 R^1 and R^2 each independently are hydrogen; C_{1-6} alkyl; C_{1-6} alkylcarbonyl; trihalomethylcarbonyl; C_{1-6} alkyl substituted with hydroxy, C_{1-6} alkyloxy, carboxyl, C_{1-6} alkylcarbonyloxy, C_{1-6} alkyloxycarbonyl or aryl; or R^1 and R^2 taken together with the nitrogen atom to which they are attached may form a morpholinyl ring or a radical of formula :



15



(d)



(e)


wherein :


R^9 , R^{10} , R^{11} and R^{12} each independently are hydrogen, halo, trihalomethyl, or C_{1-6} alkyl;

20 m is zero, 1, 2, or 3;


R^{13} , R^{14} , R^{15} and R^{16} each independently are hydrogen or C_{1-6} alkyl; or

R^{15} and R^{16} taken together may form a bivalent radical C_{4-5} alkanediyl;

- R^{17} is hydrogen; C_{1-6} alkyl; C_{1-6} alkylcarbonyl; trihalomethylcarbonyl;
 C_{1-6} alkyloxy; aryl; di(aryl)methyl; C_{1-6} alkyl substituted with hydroxy,
 C_{1-6} alkyloxy, carboxyl, C_{1-6} alkylcarbonyloxy, C_{1-6} alkyloxy; aryl;
 each R^3 independently is halo, cyano, hydroxy, trihalomethyl, trihalomethoxy,
 5 carboxyl, nitro, amino, mono- or di(C_{1-6} alkyl)amino, C_{1-6} alkylcarbonylamino,
 aminosulfonyl, mono- or di(C_{1-6} alkyl)aminosulfonyl, C_{1-6} alkyl, C_{1-6} alkyloxy,
 C_{1-6} alkylcarbonyl or C_{1-6} alkyloxy; aryl;
 each R^4 independently is halo, cyano, hydroxy, trihalomethyl, trihalomethoxy,
 carboxyl, nitro, amino, mono- or di(C_{1-6} alkyl)amino, C_{1-6} alkylcarbonylamino,
 10 aminosulfonyl, mono- or di(C_{1-6} alkyl)aminosulfonyl, C_{1-6} alkyl, C_{1-6} alkyloxy,
 C_{1-6} alkylcarbonyl or C_{1-6} alkyloxy; aryl;
 each R^5 independently is C_{1-6} alkyl, cyano or trihalomethyl;
 X is CR^6R^7 , NR^8 , O, S, $S(=O)$ or $S(=O)_2$; wherein
 R^6 and R^7 each independently are hydrogen, hydroxy, C_{1-6} alkyl, trihalomethyl,
 15 C_{1-6} alkyloxy or R^6 and R^7 taken together may form methylene; mono- or
 di(cyano)methylene; a bivalent radical of formula $-(CH_2)_2-$, $-(CH_2)_3-$, $-(CH_2)_4-$, $-(CH_2)_5-$,
 $-O-(CH_2)_2-O-$, $-O-(CH_2)_3-O-$; or, together with the carbon atom to which
 they are attached, a carbonyl;
 R^8 is hydrogen, C_{1-6} alkyl, C_{1-6} alkylcarbonyl, arylcarbonyl, aryl C_{1-6} alkylcarbonyl,
 20 C_{1-6} alkylsulfonyl, arylsulfonyl or aryl C_{1-6} alkylsulfonyl;
 each  independently represents a bivalent aromatic heterocycle wherein the
 heterocycle may be selected from the group consisting of pyrrole, pyrazole, imidazole,
 triazole, furane, thiophene, isoxazole, oxazole, isothiazole, thiazole, pyridine,
 pyridazine, pyrimidine, pyrazine and triazine; or one of the two bivalent aromatic
 25 heterocycles in the compounds of formula (I) may be 1,2-benzenediyl; and
 aryl is phenyl; or phenyl substituted with 1, 2 or 3 substituents selected from halo,
 hydroxy, C_{1-6} alkyl and trihalomethyl.


2. A compound according to claim 1 wherein one of the two  radicals is
 1,2-benzenediyl.
 30 3. A compound according to claim 1 or 2 wherein X is CR^6R^7 or O.
 4. A compound according to any one of claims 1 to 3 wherein R^1 and R^2 are both
 methyl and n is 1 or 2.
 35


5. A compound as claimed in claim 1 wherein the compounds is
 2,3,3a,8-tetrahydro-*N,N*-dimethylisoxazolo[2,3-*a*]pyrrolo[2,1-*c*][1,4]benzo-
 diazepine-2-methanamine;
 2,3,3a,8-tetrahydro-*N,N*-dimethylimidazo[2,1-*c*]isoxazolo[2,3-*a*][1,4]benzo-
 diazepine-2-methanamine;
 2,3,3a,7-tetrahydro-*N,N*-dimethylisoxazolo[2,3-*a*]thieno[2,3-*c*][1]benz-azepine-2-
 methanamine;
 2,3,3a,7-tetrahydro-*N,N*-dimethylisoxazolo[2,3-*a*]thieno[3,2-*c*][1]benz-azepine-2-
 methanamine; a stereochemically isomeric form or a pharmaceutically acceptable
 addition salt thereof, or an *N*-oxide form thereof.
6. A composition comprising a pharmaceutically acceptable carrier and, as active
 ingredient, a therapeutically effective amount of a compound as claimed in any one
 of claims 1 to 5.
7. A process of preparing a composition as claimed in claim 6, wherein a
 pharmaceutically acceptable carrier is intimately mixed with a therapeutically
 effective amount of a compound as claimed in any one of claims 1 to 5.
8. A compound as claimed in any one of claims 1 to 5 for use as a medicine.
9. A compound of formula (II)



wherein , X, R³, R⁴, p and q are as defined in claim 1.

10. A process of preparing a compound as described in claim 1, characterized in that :
- a) a dienophile of formula (III) is reacted with an intermediate of formula (II) :



wherein in the intermediates (II) and (III) R^1 to R^5 , X , n , r , p , q and  are

defined as in claim 1, and the hydrogens on the sp^2 -hybridized aliphatic carbon atoms may each individually be replaced by R^5 ;

- 5 b) converting compounds of formula (I) into each other following art-known transformations, and further, if desired, converting the compounds of formula (I), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or into a therapeutically active non-toxic base addition salt by treatment with a base, or
10 conversely, converting the acid addition salt form into the free base by treatment with alkali, or converting the base addition salt into the free acid by treatment with acid;
and, if desired, preparing stereochemically isomeric forms or *N*-oxide forms thereof.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 97/01830

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D498/16 C07D495/04 C07D487/04 C07D471/04 C07D498/04
A61K31/55 //(C07D498/16,333:00,261:00,223:00),(C07D498/16,
261:00,243:00,209:00),(C07D495/04,333:00,223:00),(C07D487/04,

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	WO 96 14320 A (JANSSEN PHARMACEUTICA N.V.) 17 May 1996 see claims 1,8-11,13 ---	1,6-10
A,P	WO 96 14321 A (JANSSEN PHARMACEUTICA N.V.) 17 May 1996 see claims 1,6-9,11 ---	1,6-10
A	US 4 039 558 A (W. J. VAN DER BURG ET AL.) 2 August 1977 cited in the application see column 3, line 53 - line 65; claim 1 ---	1,6,8
A	EP 0 421 823 A (SANKYO CO. LTD.) 10 April 1991 cited in the application see claims 1,40,42 ---	1,6,8
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

29 July 1997

Date of mailing of the international search report

0 8. 08. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Hass, C

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 97/01830

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 243:00,209:00), (C07D471/04,223:00,221:00), (C07D498/04,267:00,221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JOURNAL OF HETEROCYCLIC CHEMISTRY, vol. 17, no. 2, 1980, pages 341-50, XP002036279 R. B. MOFFETT: "Tetracyclic Heterocycles as CNS Agents" see table I ---	1
A	TETRAHEDRON, vol. 46, no. 20, 1990, pages 7207-18, XP002036280 S. A. ALI ET AL.: "The 1,3-dipolar cycloaddition Reactions of 3,4,5,6-tetrahydro-2H-azepine 1-oxide" --- -/--	10



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

08.08.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 97/01830

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	TETRAHEDRON LETTERS , vol. 28, no. 3, 1987, pages 263-6, XP002036281 J. J. TUFARIELLO ET AL.: "A Highly Stereoselective Synthesis of (E,E)-1,5-Dienes" see page 263, reaction scheme ---	10
A	TETRAHEDRON LETTERS, vol. 28, no. 3, 1987, pages 267-70, XP002026036 J.J. TUFARIELLO ET AL.: "The Synthesis and Cycloaddition Reactions of 3-Azabicyclo[3.1.0]hex-2-ene 3-oxide and 3-Azabicyclo[3.2.0]hept-2-ene 3-oxide. Highly strained Bicyclic Nitrones" see table 1 ---	10
A	TETRAHEDRON LETTERS, vol. 28, no. 23, 1987, pages 2649-52, XP002036282 G. PANDEY ET AL.: "Photosensitized single electron transfer oxidation of N-hydroxylamines: A convenient synthesis of cyclic nitrones" see table I -----	10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/01830

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9614320 A	17-05-96	AU 3924995 A FI 971854 A NO 972017 A US 5552399 A	31-05-96 30-04-97 30-04-97 03-09-96
WO 9614321 A	17-05-96	AU 3925095 A FI 971855 A NO 972018 A	31-05-96 30-04-97 30-04-97
US 4039558 A	02-08-77	NL 7414038 A BE 834902 A CA 1082182 A DE 2548045 A FR 2289198 A GB 1530299 A JP 51068598 A SE 7511968 A ZA 7506610 A	03-05-76 27-04-76 22-07-80 06-05-76 28-05-76 25-10-78 14-06-76 29-04-76 29-09-76
EP 421823 A	10-04-91	CA 2026925 A CN 1051360 A,B EP 0505014 A JP 3279383 A JP 8019130 B NO 175781 B RU 2048469 C US 5476848 A US 5461051 A US 5362725 A	06-04-91 15-05-91 23-09-92 10-12-91 28-02-96 29-08-94 20-11-95 19-12-95 24-10-95 08-11-94